

الملح الصخري في اليمن*

()

(Na CL)

-1

-2

-3

% 78

³ / 2.2

2.5-2

.

. 800°

-:

.

.

200

1991

.

31.4

()

()

)

.(

-1

100 - 40

)

-(

-1-1

80

1972 ()

33

95

31.23

% 98

% 34.94	
% 0.016	
% 0.36	
% 0.0083	
% 59.60	
% 0.075	
% 0.38	
% 98.23	

(1)

. 80

115

.2

37

2 2

. 150

()

.3

97

% 98 - 96

.4

()

()

- 1-4

4

47

. 35 - 30

² 4

- 2-4

15 86
25 - 20 2 1

- 3-4

120 18
16-14 2 1

-4-4

20

()

10 - 7

.2 2

%								
Fe	K	Mg	Ca	So ₄		Nacl		
0.028	0.16	0.036	0.1	0.33	4.40	94.35		Nacl 2
0.017	0.12	0.049	0.16	0.45	1.80	96.86		Nacl 3
0.11	0.24	0.061	0.5	1.98	5.80	91.48		Nacl 4
0.11	0.11	0.097	3.61	7.41	29.40	54.43		Nacl 5

(2)

1952

150

410

1984

/ 7

18

18.5

-1

-2

92	91	90	89	88	87	86	85	84	
107	115	143	152	135	135	135	135	135) (
95	95	120	120	120	120	120	120	120	()
15	20	23	32	15	15	15	15	15	()

(3)

. / 8 / 850

. 410

-1

-2

. 150,000

. 8.5

-3

-4

. 410 1997

-5

-6

-7

-8

-1

-2

-3

-4

..... (1- 1)

..... (1)

..... (2)

..... (3)

..... (1)

..... (2)

2000

. 1997 - 94

. 1993 - 84

. 1972

-

-

-1

-2

-3

-4

-5

1. Geomin 1982.

- 2. H.N,Mills, Industrial Minerals and Rocks (Vol.2).**
- 3. Strojex port Prague – Czech – Final Report on Economic Geology of Non- metallic Raw material in Yemen 1977.**